



<b>Abbreviated name</b>	Condom use at last sex with high-risk partner
<b>Indicator name</b>	Condom use at last sex with high-risk partner
<b>Domain</b>	Risk factors
<b>Subdomain</b>	Infectious disease
<b>Associated terms</b>	Infections
<b>Definition</b>	Percentage using a condom during last sexual intercourse with a higher-risk partner (women and men who have more than one sexual partner in the past 12 months; sex workers with most recent client; men who have sex with men anal sex with a male partner; people who inject drugs condom use at last sex). For people who inject drugs, also measure number of needles per person who injects drugs per year.
<b>Numerator</b>	
<b>Denominator</b>	
<b>Disaggregation/ additional dimension</b>	Age (15–24, 15–49 years), sex Sex workers: by sex (F/M/transgender) age (< 25/25+ years) Men who have sex with men: age (< 25/25+ years) People who inject drugs: sex, age (< 25/25+ years)
<b>Method of measurement</b>	Population-based surveys for general population; Surveys targeting key populations such as IBBS.
<b>Method of estimation</b>	
<b>Measurement frequency</b>	
<b>Monitoring and evaluation framework</b>	Outcome
<b>Preferred data sources</b>	Surveys
<b>Other possible data sources</b>	
<b>Further information and related links</b>	The Demographic and Health Surveys Program. Washington (DC): United States Agency for International Development ( <a href="http://www.dhsprogram.com/">http://www.dhsprogram.com/</a> , accessed 29 March 2015).