



Intermittent preventive therapy for malaria during pregnancy (IPTp)

Abbreviated name	Intermittent preventive therapy for malaria during pregnancy (IPTp)
Indicator name	Intermittent preventive therapy for malaria during pregnancy (IPTp)
Domain	Service coverage
Subdomain	Infectious disease
Associated terms	Malaria
Definition	Percentage of women who received three or more doses of intermittent preventive treatment during antenatal care visits during their last pregnancy.
Numerator	Number of women receiving three or more doses of recommended treatment.
Denominator	Total number of pregnant women/surveyed with a live birth in the last 2 years.
Disaggregation/ additional dimension	Age, place of residence, socioeconomic status
Method of measurement	
Method of estimation	
Measurement frequency	
Monitoring and evaluation framework	Outcome
Preferred data sources	Household surveys, facility information systems
Other possible data sources	
Further information and related links	<p>Countdown to 2015 decade report (2000–2010): taking stock of maternal, newborn and child survival. Geneva and New York (NY): World Health Organization/United Nations Children's Fund; 2010 (http://www.countdown2015mnch.org/reports-and-articles/previous-reports/2010-decade-report, accessed 29 March 2014).</p> <p>Household Survey Indicators for Malaria Control. Measure Evaluation/Measure DHS/President's Malaria Initiative/Roll Back Malaria Partnership/UNICEF/WHO. 2013 (http://www.malariasurveys.org/documents/Household%20Survey%20Indicators%20for%20Malaria%20Control.pdf, accessed 15 April 2015).</p> <p>Roll Back Malaria Partnership/WHO. Disease surveillance for malaria control: an operations manual. Geneva: World Health Organization; 2012 (http://www.who.int/malaria/publications/atoz/9789241503341/en/, accessed 29 March 2015).</p> <p>World Health Assembly governing body documentation: official records. Geneva: World Health Organization (http://apps.who.int/gb/or/, accessed 29 March 2015).</p>