


Analysis and use
of community-based
health service data

GUIDANCE FOR COMMUNITY HEALTH WORKERS STRATEGIC INFORMATION AND SERVICE MONITORING



SELECTED HIGHLIGHTS



Published by UNICEF
Division of Communication
3 United Nations Plaza
New York, NY 10017, USA

Contact: pubdoc@unicef.org

Website: www.unicef.org

Suggested citation:

United Nations Children's Fund, World Health Organization,
Global Fund to Fight AIDS, Tuberculosis and Malaria,
United Nations Entity for Gender Equality and the
Empowerment of Women, and Gavi, the Vaccine Alliance,
*Guidance for community health worker strategic information
and service monitoring: Selected highlights*, UNICEF, New York, 2021

ISBN: 978-92-806-5208-6

© United Nations Children's Fund (UNICEF), JANUARY 2021



Analysis and use
of community-based
health service data

GUIDANCE FOR COMMUNITY HEALTH WORKERS STRATEGIC INFORMATION AND SERVICE MONITORING

SELECTED HIGHLIGHTS





CONTENTS

EXECUTIVE SUMMARY	v
WHO ARE COMMUNITY HEALTH WORKERS	1
SCOPE	3
ALIGNMENT WITH EXISTING MONITORING FRAMEWORKS	4
MAIN STEPS TO CONSIDER WHEN CHOOSING MODULES AND INDICATORS	6
1 Review national strategies to prioritize modules	6
2 Review CHW tasks and CHIs maturity by indicator	6
3 Review the reporting burden of CHWs	6
4 Review the quality of reporting	6
5 Review data use, equity and feedback loops	7
LIST OF INDICATORS	9

LIST OF FIGURES

Figure 1	Alignment of malaria indicators from data collection at the community level to global technical strategy	5
Figure 2	Main steps to consider when choosing modules and indicators	7



EXECUTIVE SUMMARY

Community health workers (CHWs) are non-professional health workers, either paid or volunteer, who are based in communities and provide outreach beyond primary health-care facilities or who are based at health posts not staffed by professional medical staff. Compared to other types of health workers, CHW cadres across and even within countries are remarkably diverse in terms of their tasks, functions and degree of institutionalization into the formal health sector.

A major obstacle to delivering services of good quality to children, adolescents and adults – including to key and vulnerable populations at the community level – is the dearth of data on community-level care and how they are managed and used. The absence of standardized and aligned CHW services indicators still limits the development of well-functioning community health information systems (CHIS), their integration within broader health information systems (HIS) and monitoring and evaluation (M&E) frameworks inclusive of both community- and facility-based service data. To facilitate the harmonization of monitoring of CHW services, this guidance offers a modular set of common indicators reported by CHWs at the time they provide services.

These indicators are aligned with existing monitoring frameworks, and are flexible enough to be adapted for different country contexts and varying maturity levels of different CHIS.

Areas covered by this guidance are:

- **population composition**
- **water, sanitation and hygiene (WASH)**
- **clean energy**
- **sexual and reproductive health (SRH)**
- **maternal health**
- **newborn health**
- **child health**
- **adolescent health**
- **immunization**
- **HIV**
- **malaria**
- **tuberculosis (TB)**
- **neglected tropical diseases (NTDs)**
- **child protection and interpersonal violence**
- **civil registration and vital statistics (CRVS)**
- **non-communicable diseases (NCDs)**
- **nutrition**
- **mental health**
- **people-centred services**
- **community-based surveillance (CBS)/early warning.**

Annex 1 of the Main Document and Indicators report lists the indicators by module, and a separate volume accompanying this guidance details the metadata for each indicator.

During the process of selection of indicators, six principles are critical for a human rights–based approach to data: participation, data disaggregation, self-identification, transparency, privacy and accountability. It is important to ensure and plan for the participation of all relevant stakeholders, including for planning, data collection, dissemination and analysis of data. The views of vulnerable or marginalized groups, and groups who are at risk of discrimination, should be represented. These principles are critical in designing an equity-focused process, well articulated with community-led monitoring, to address specific data use cases, and inform all stakeholders for targeted action to reach the 2030 Sustainable Development Agenda.

The following are the key steps to choosing and standardizing CHW indicators for the main implementing partners:

1. REVIEW NATIONAL STRATEGIES TO PRIORITIZE MODULES



COUNTRY LEADERS

- Identify who is responsible for setting up or updating the CHIS, especially in terms of the standardization of CHW indicators.
- Map key actors, including communities, the private sector and civil society organizations, implementers (nongovernmental organization [NGOs], including for community-based surveillance [CBS]) and all other key stakeholders.
- Review all relevant national or subnational strategies.
- Evaluate carefully the plans and ongoing initiatives for digitalization and their impact in terms of the choice of indicators, including for maturity and costs.
- Review the two main types of CHW activities (consultations versus household assessments), including which tasks CHWs are allowed to perform.
- Identify the specific CHW indicator modules to prioritize.

IMPLEMENTING PARTNERS, NGOs and PRIVATE SECTOR

- Actively participate in the process and provide feedback on key modules to focus on.
- Plan how best to align and standardize CHW indicators, and how best to contribute to CHIS.

COMMUNITIES/CIVIL SOCIETY ORGANIZATIONS

- Actively participate in the process and provide feedback on key modules to focus on.
- Allow community-led organisations to bring their experience in community led monitoring to inform CHW strategic information.

FUNDING PARTNERS

- Review with country leaders and implementing partners costs and sustainability of the plan.
- Ensure good governance of the whole process of standardization of CHW indicators.

ACADEMIA

- Review whether there are opportunities to learn from the standardization and digitalization process to facilitate it – for instance, through implementation research.

2. REVIEW CHW TASKS AND CHIS MATURITY BY INDICATOR



COUNTRY LEADERS

- For each chosen module, review which CHW tasks should be reported during household assessments or CHW consultation types of activities.
- Choose CHW indicators based on CHW tasks as a primary filter.
- Then, review CHW indicators based on their specific maturity within the CHIS.

IMPLEMENTING PARTNERS, NGOs and PRIVATE SECTOR

- Review which tasks CHWs are doing and coordinate with country leaders so these are captured by CHIS.

3. REVIEW THE REPORTING BURDEN OF CHWs



COUNTRY LEADERS

- Estimate the CHW reporting burden by considering the total number of data points to be collected, the number of indicators to calculate, disaggregation, frequency of reporting, the design of data collection instruments and the average catchment area.
- Based on this assessment, review the number of indicators, their disaggregation and reporting frequency; prioritize indicators and basic types of disaggregation.

IMPLEMENTING PARTNERS, NGOs and PRIVATE SECTOR

- Actively participate in the selection process and provide feedback on key CHW tasks and corresponding indicators to focus on, as well as CHW reporting burden.
- Discuss alignment in terms of frequency of reporting and disaggregation.
- Share best practices on reporting tools.

COMMUNITIES/CIVIL SOCIETY ORGANIZATIONS

- Actively participate in the process and provide feedback on key indicators to focus on.
- Contribute to and articulate the choice of indicators with community-led monitoring.

FUNDING PARTNERS

- Review with country leaders and implementing partners costs and sustainability of the tools, especially if undergoing digitalization.

ACADEMIA

- Review the best ways to estimate the reporting burden of CHWs.



© UNICEF/UNI106359/Pirozzi

4. REVIEW THE QUALITY OF REPORTING



COUNTRY LEADERS

- For each indicator considered, assess potential double counting and how it can be minimized.
- Explore the feasibility of clinical assessment indicators in terms of resources, training and reliability.
- Consider all levels of CHIS interoperability and whether they are taken into account within a comprehensive health information system strategy.
- Ensure necessary policies or legal framework and standard operating procedures (SOPs) are in place to ensure confidentiality.
- Review whether there are issues with processes and practices ensuring confidentiality.
- Review whether data security in all its dimensions can be ensured.
- Based on these evaluations and a review of data quality in general, consider whether certain CHW indicators should be measured or tested for feasibility before scaling up.

IMPLEMENTING PARTNERS, NGOs and PRIVATE SECTOR

- Offer feedback on issues and best practices on data quality, feasibility, confidentiality and data security for the indicators considered.
- Review specifically issues of double counting between implementing partners.
- Review how best to operationalize interoperability to best contribute to CHIS.
- Discuss alignment in terms of measurement, including for clinical assessments.

COMMUNITIES/CIVIL SOCIETY ORGANIZATIONS

- Provide feedback on issues of data quality, confidentiality and data security for the indicators considered.
- Offer possible solutions or share known best practices.

FUNDING PARTNERS

- Review how best to support improvement initiatives for data quality with data use, interoperability and data security/confidentiality.
- Review how to support the sustainability of training, supervision and reporting resources.

ACADEMIA

- Research on best ways to improve data quality, interoperability and data security/confidentiality.

5. KEY STEPS FOR ACTION WHEN REVIEWING DATA USE, EQUITY AND FEEDBACK LOOPS



COUNTRY LEADERS

- Aim for a rights-based strategy for data democratization.
- Review whether there are multicomponent interventions in place to improve data quality, access and use.
- Establish clear SOPs for each chosen indicator, including the ways it should be collected, calculated (with denominators if relevant), analysed and provided feedback on, and possible actions it could trigger.
- Ensure communities are included in the processes with feedback loops, convening participatory routine data reviews with community/civil society partners.
- Establish the monitoring approach for equity in general (including gender, age and geographic location) and vulnerable and key populations in particular.

IMPLEMENTING PARTNERS, NGOs and PRIVATE SECTOR

- Articulate different feedback loops between partners.
- Review and articulate equity strategies and their monitoring between implementing partners, including for vulnerable populations.

COMMUNITIES/CIVIL SOCIETY ORGANIZATIONS

- Put in place feedback monitoring with country leaders and implementing partners.
- Articulate CHW indicators with community-led monitoring.
- Participate in assessing inequities, underserved populations, and human rights and gender related barriers to services.
- Representatives of vulnerable populations should supervise equity monitoring strategy.


FUNDING PARTNERS

- Ensure good governance in terms of data use, feedback loops and equity strategy.

ACADEMIA

- Research on best ways to improve feedback loops, equity, articulation of different community monitoring systems, including community-led monitoring, and monitoring and participation of vulnerable populations.

Je donne uniquement mon lait à mon bébé
dès la naissance jusqu'à 6 mois



MINISTÈRE DE LA SANTÉ
BURKINA FASO



unicef







WHO ARE COMMUNITY HEALTH WORKERS

Community health workers (CHWs) are non-professional health workers, either paid or volunteer, who are based in communities and provide outreach beyond primary health-care facilities or who are based at health posts not staffed by professional medical staff. Compared to other types of health workers, CHW cadres across and even within countries are remarkably diverse in terms of their tasks, functions and degree of institutionalization into the formal health sector.

In many health systems, these activities fall in a continuum between two main types of activities: individual consultations and household assessments.

- **Ongoing individual consultations:** These are consultations performed by CHWs on a regular basis, and can either happen at the health post or during home visits. This constitutes the core of their activities.
- **Household assessments:** In certain countries, CHWs do annual or semi-annual household assessments that are akin to survey or census activities. These visits usually aim to assess some characteristics of the population, of specific practices (vector control, WASH), but also to deliver some interventions.



SCOPE

Due to the fact that **CHWs are charged with diverse tasks in different countries, this guidance offers a modular set of common indicators reported by CHWs at the time they provide services.** This format offers flexibility to fit the national community health strategies, the epidemiological situation and the key functions of CHWs in specific countries (for example, indicators on malaria will not be needed in non-malaria settings). This document is a derivative guidance based on multiple guidelines. The proposed set of community indicators has been reviewed so their metadata align as much as possible with published guidelines, and to allow the data to flow from the community level to the facilities, district, national and global levels, where necessary.

Areas covered by this guidance are:

- **population composition**
- **water, sanitation and hygiene (WASH)**
- **clean energy**
- **sexual and reproductive health (SRH)**
- **maternal health**
- **newborn health**
- **child health**
- **adolescent health**
- **immunization**
- **HIV**
- **malaria**
- **tuberculosis (TB)**
- **neglected tropical diseases (NTDs)**
- **child protection and interpersonal violence**
- **civil registration and vital statistics (CRVS)**
- **noncommunicable diseases (NCDs)**
- **nutrition**
- **mental health**
- **people-centred services**
- **community-based surveillance (CBS)/ early warning.**



© UNICEF/UNI45453/Pirozzi

ALIGNMENT WITH EXISTING MONITORING FRAMEWORKS


To facilitate data flow, data aggregation and interoperability across the HIS, indicators should be aligned from the community level to the global level as much as possible. Typically, global strategies and frameworks – such as the Monitoring Framework for the Sustainable Development Goals or UN Every Woman Every Child Global Strategy – have been endorsed by countries after comprehensive processes of consultation and validation. WHO publishes normative guidelines as global goods; for example, the WHO Consolidated HIV strategic information guidelines are based on a robustly representative, participatory/ partner-based process.

This derivative guidance is based on multiple guidelines. The proposed set of community indicators has been reviewed so their metadata align as much as possible with published guidelines, and to allow the data to flow from the community level to the facilities, district, national and global levels, where necessary. The figure below depicts the data flow from the community to global strategic planning for malaria.



© UNICEF/UN0267026/Raoulison

Figure 1 Alignment of malaria indicators from data collection at the community level to global technical strategy



ACT: artemisinin-based combination therapy; IPTp: intermittent preventive treatment of malaria during pregnancy; ITN: insecticide-treated nets; SMC: seasonal malaria chemoprevention

MAIN STEPS TO CONSIDER WHEN CHOOSING MODULES AND INDICATORS

1. Review national strategies to prioritize modules

To facilitate the integration of CHIS within the HIS, adopting a health systems perspective is particularly critical. As the community health system must be integrated into the overall national health system, the CHIS must also be integrated with the national HIS, ensuring alignment with existing national strategies, policies and action plans. These national strategies will inform which modules should be prioritized in developing CHW indicators. Sustainable resources available for the CHIS should also be estimated and budgeted properly. Digitalization can become a phenomenal accelerator if carefully designed with a health system perspective and bringing all stakeholders to the table from the start, including end-users.

2. Review CHW tasks and CHIS maturity by indicator

The list of indicators in this guidance consist of a menu from which countries can select desired indicators; countries should not use all indicators. For each module, CHW tasks are listed – corresponding indicators should be considered only if these tasks are performed or planned in the country. The same indicators can appear in different modules as they may be critical for different modules (for example, child health and nutrition). Supervisors and programme managers must keep in mind that not all CHW key activities may be reflected in the routine CHIS. Because the development and design of CHIS are frequently service specific, CHWs usually have different ways of reporting information, depending on their tasks, with different registers for different programmes. Therefore, a simplified maturity score has been assigned to each indicator and not to the whole CHIS. For instance, longitudinal tracking systems with unique IDs at CHW level may be available only for follow-up of pregnancies. A given indicator should be considered then only if the CHIS has the maturity necessary for this specific indicator to be reported.

3. Review the reporting burden of CHWs

After considering the different pieces of community monitoring, and the best way to articulate them given needs, resources and capabilities, the CHW workload and reporting burden are critical to map and evaluate as they can impact the quality of the data reported. The indicators presented in this document should be considered part of the whole reporting burden for CHWs. Deciding how, by whom and at what level the indicator should be calculated is therefore a critical decision when designing the CHIS. Frequency of reporting can vary a lot between indicators, according to the objective of the indicator and the specific country HIS. Based on this review of burden, prioritizing and reducing the number of indicators may become a necessity. For each module, indicators have been classified as either priority or additional, to help with the prioritization exercise. Countries might first consider the priority indicators if they need to reduce the reporting burden.

4. Review the quality of reporting


The quality of CHW services and reporting should be assessed when choosing which indicators can be reported, and then periodically reassessed. Critical elements to review are the design of reporting tools, coordination and adequate training of all people involved in the CHIS: collection, cleaning, analysis, use and communication of data. Some key elements in particular are to be checked when selecting indicators: unintentional double or multiple counting of the same event, accuracy of indicators reporting on clinical assessments such as anthropometric measurements, interoperability, confidentiality and data security processes.

5. Review data use, equity and feedback loops

Data collection should be used to inform decision-making for action to improve programmes. This concerns all stakeholders, and data should be fed back to the community and used by all relevant CHIS stakeholders. Data democratization can be enabled by a supporting comprehensive data use culture. Feedback of data (data loops) to all stakeholders has shown its potential to improve surveillance systems as well as the quality of health services and the corresponding data.

Targeting equity is paramount to reach the SDG agenda of “leaving no one behind and reaching the furthest behind first.” Gender-responsive health policies, programmes, services and delivery models require considering power dynamics within communities and between individuals and the differential needs of men, women, boys, girls and gender-nonconforming people in all their diversity. Vulnerable populations are essential to consider for equity, as they have a heightened risk of poor health, reduced access to services, and can be criminalized or marginalized. Characteristics of vulnerable populations need to be explicitly integrated in monitoring frameworks, so they are not left out of reporting.

Figure 2 Main steps to consider when choosing modules and indicators



CHIS: community health information system; **CHW:** community health worker; **CRVS:** civil registration and vital statistics; **IPV:** interpersonal violence; **NCDs:** noncommunicable diseases; **NTDs:** neglected tropical diseases; **TB:** tuberculosis; **WASH:** water, sanitation and hygiene



LIST OF INDICATORS

HH: household assessment; **CS:** individual consultation

The same indicators can appear in different modules as they may be critical for different modules (for example, child health and nutrition).

For each module, indicators have been classified as either **priority** (in **bold**) or additional, to help with the prioritization exercise.

Population composition

1	Number of households in the catchment area	HH
2	Number of live births	HH
3	Number of infants in the catchment area (0 to less than 1 year old)	HH
4	Number of children in the catchment area (1 to less than 5 years old)	HH
5	Number of children in the catchment area (5 to less than 10 years old)	HH
6	Number of young adolescents in the catchment area (10–14 years old)	HH
7	Number of older adolescents in the catchment area (15–19 years old)	HH
8	Number of pregnant women in the catchment area	HH
9	Number of adults in the catchment area	HH

Water, sanitation and hygiene (WASH)

10	Main drinking-water source	HH
11	Time to collect drinking-water	HH
12	Location of drinking-water point	HH
13	Drinking-water quality at the source	HH
14	Availability of drinking-water when needed	HH
15	Access to handwashing facilities with water and soap available on premises (household level)	HH
16	Use of improved sanitation facilities	HH
17	Sharing of sanitation facilities	HH
18	Emptying of on-site sanitation facilities (septic tanks and pit latrines)	HH
19	Private place to wash and change during menstruation	HH
20	Use of hygiene materials during menstruation	HH

Clean energy

21	Proportion of households with primary reliance on clean fuels and technologies for cooking	HH
22	Proportion of households with primary reliance on clean fuels and technologies for lighting	HH
23	Proportion of households with primary reliance on clean fuels and technologies for heating	HH

Sexual and reproductive health

24	Number of condoms distributed	HH, CS
25	Number of women and adolescent girls who initiate modern methods for family planning in the community	CS
26	Number of women and adolescent girls who use modern methods for family planning	HH
27	Proportion of adolescents and young people seeking contraception/family planning who received an HIV test	CS
28	Number of HIV tests conducted (testing volume) and the proportion of HIV-positive results returned to people (positivity)	HH, CS
29	Men and adolescent boys with urethral discharge	CS
30	Proportion of antenatal care attendees tested for syphilis	CS
31	Proportion of women who have been screened for cervical cancer	HH
32	Number of women referred for any post-abortion complication	CS
33	Number of women referred for safe abortion	CS
34	Number/proportion of women aged 15–49 years old who have undergone female genital mutilation	HH

Maternal health

35	Proportion of pregnant women who have first antenatal care (ANC) contact with CHW in first trimester	CS
36	Proportion of antenatal care contacts during which women received breastfeeding counselling	CS
37	Proportion of community antenatal care contacts in the reporting period during which pregnant women were given/prescribed iron-containing supplements	CS
30	Proportion of antenatal care attendees tested for syphilis	CS
38	Proportion of people with raised blood pressure measured by CHW	HH, CS
28	Number of HIV tests conducted (testing volume) and the proportion of HIV-positive results returned to people (positivity)	HH, CS
39	Number of pregnant women referred for maternal complications	CS
40	Proportion of women who gave birth in the community without skilled birth attendant and who were administered oral immediate postpartum uterotonic to prevent postpartum haemorrhage	CS
41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH
44	Number of pregnancy-related deaths	HH

Newborn health

45	Number of stillbirths (late fetal deaths)	HH
46	Number of neonatal deaths (0–27 days)	HH
47	Proportion of newborns delivered in the community who were put to breast within the first hour of birth	CS
48	Proportion of newborns delivered in the community with documented birthweight	CS
49	Prevalence of low birthweight among newborns delivered in the community	CS
50	Number of preterm newborns discharged from facility that received follow-up on Kangaroo Mother Care (KMC) by CHW	CS

51	Proportion of preterm newborns delivered in the community	CS
52	Number/proportion of newborns and children referred for danger signs	CS
53	Proportion of newborns delivered in the community initiated on skin-to-skin contact immediately after birth	CS
54	Newborns delivered in the community whose cord was cut with clean instrument	CS

Malaria

113	Number/proportion of children under 5 who had fever in the last two weeks	HH
114	Number/proportion of households with at least one insecticide-treated nets (ITN)	HH
115	Number/proportion of households with at least one ITN for every two persons	HH
116	Number/proportion of population who slept under an ITN the previous night	HH
117	Number/proportion of population living in a house sprayed by IRS in the previous 12 months	HH
118	Number of ITNs distributed by CHWs	HH, CS
119	Number/proportion of febrile cases who received a rapid diagnostic test (RDT)	CS
120	Number/proportion of suspected cases who received a RDT	CS
121	Malaria RDT positivity rate	CS
122	Number/proportion of patients with malaria who received first-line antimalarial treatment according to national policy	CS
123	Number/proportion of children with confirmed malaria and danger signs who are referred	CS
124	Number/proportion of children with malaria danger signs referred after administration of rectal artesunate	CS
125	Number of pregnant women who have received zero/one/two/three or more doses of intermittent preventive treatment of malaria in pregnancy (IPTp)	CS
126	Number of children aged 3–59 months who received zero/one/two/three/four or more courses of seasonal malaria chemoprevention (SMC) in a transmission season	HH, CS

Child health

55	Number of infant deaths (0 to less than 1 year old)	HH
56	Number of child deaths (1 to less than 5 years old)	HH
57	Number of child deaths (5 to less than 10 years old)	HH
58	Proportion of consultations for infants under 6 months providing counselling on appropriate infant and young child feeding	CS
59	Proportion of consultations for children 6–23 months providing counselling on appropriate complementary feeding	CS
60	Proportion of people receiving preventive chemotherapy for deworming	CS
61	Percentage of children 6–59 months of age who received an age-appropriate dose of vitamin A through CHW contacts (routine contacts as well as contacts via events) in each semester	CS
62	Number/proportion of children aged 6–59 months assessed for wasting in the reporting period	CS
63	Proportion of children 6–59 months with mid-upper-arm circumference (MUAC) < 115 mm (severe acute malnutrition)	CS
64	Number/proportion of children under 5 years who had their weight assessed	CS
65	Number/proportion of children under 5 years who are underweight	HH, CS
66	Proportion of wasting among children aged 6–59 months	CS
67	Number/proportion of children under 5 years who had their height/length measured	CS
68	Proportion of overweight among children under 5 years of age	HH, CS
69	Proportion of overweight and obesity in school-age children and adolescents 5–19 years	HH, CS
70	Number/proportion of children who received treatment for diarrhoea	CS

71	Proportion of children presenting fast breathing and/or chest indrawing	CS
72	Proportion of children receiving antibiotic treatment for fast breathing and/or chest indrawing	CS
73	Number/proportion of young infants, 0–59 days old who received pre-referral treatment for signs of possible serious bacterial infection	CS
74	Number/proportion of caregivers receiving information on early identification of danger signs	CS
52	Number/proportion of newborns and children referred for danger signs	CS
75	Proportion of children monitored for early signs of developmental delays	CS
76	Proportion of children with suspected developmental disabilities referred	CS
77	Proportion of caregivers being counselled on responsive caregiving and early learning activities	CS
78	Proportion of children with suspected developmental disorders and disabilities whose caregivers receive parenting information	CS
28	Number of HIV tests conducted (testing volume) and the proportion of HIV-positive results returned to people (positivity)	HH, CS
79	Number of people referred for HIV testing	HH, CS
80	Insufficient physical activity among children (0 to less than 5 years of age)	CS
81	Insufficient physical activity among children (5 to less than 10 years of age)	CS

Adolescent health

82	Number of young adolescent deaths (10–14 years old)	HH
83	Number of older adolescent deaths (15–19 years old)	HH
27	Proportion of adolescents and young people seeking contraception/family planning who received an HIV test	CS
28	Number of HIV tests conducted (testing volume) and the proportion of HIV-positive results returned to people (positivity)	HH, CS
84	Supplementation with iron-containing supplements (and folic acid) to adolescent girls and women of reproductive age for anaemia prevention	HH, CS
60	Proportion of people receiving preventive chemotherapy for deworming	CS
85	Women and adolescent girls who are underweight	HH, CS
69	Proportion of overweight and obesity in school-age children and adolescents 5–19 years	HH, CS
86	Insufficient physical activity among adolescents	HH
87	Proportion of adolescents currently using tobacco	HH, CS
88	Proportion of alcohol consumers	HH
89	Proportion of people with heavy episodic drinking	HH
90	Proportion of people who use psychoactive drugs	HH
91	Proportion of people who inject psychoactive drugs	HH
41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH
92	Number of people with suicidal ideation or plan	CS
93	Number of people with suicide attempts	CS
94	Average time on weekdays and weekend days dedicated to screen time for leisure activities	HH

Immunization

95	Number/proportion of persons who are not up to date with immunizations and are referred	HH, CS
----	--	--------

96	Number/proportion of children under 5 never vaccinated (zero-dose) and referred	HH, CS
97	Detection and reporting of neonatal tetanus	CS
98	Detection and reporting of acute flaccid paralysis	CS
99	Detection and reporting of rash and fever for measles or measles/rubella	CS
100	Polio vaccination with oral polio vaccination (OPV) (%)	CS

HIV

24	Number of condoms distributed	HH, CS
27	Proportion of adolescents and young people seeking contraception/family planning who received an HIV test	CS
28	Number of HIV tests conducted (testing volume) and the proportion of HIV-positive results returned to people (positivity)	HH, CS
79	Number of people referred for HIV testing	HH, CS
101	Early infant referral	CS
102	Number of individual HIV self-testing (HIVST) kits distributed	HH, CS
103	Number of people who were identified and tested for HIV using HIV index testing services and received their results	HH, CS
104	Proportion of people on pre-exposure prophylaxis (PrEP) supported and counselled by CHWs	CS
105	Number and percentage of people living with HIV reported on antiretroviral therapy (ART) at the end of the last reporting period and/or newly initiating ART during the current reporting period who were not on ART at the end of the reporting period	HH
106	Number of people living with HIV on ART	CS
30	Proportion of antenatal care attendees tested for syphilis	CS
31	Proportion of women who have been screened for cervical cancer	HH
107	Avoidance of health care due to stigma and discrimination (key populations)	HH
108	Avoidance of health care due to stigma and discrimination (people living with HIV)	HH
109	Proportion of people eligible for any form of economic support referred by a CHW to apply for it	CS
110	Proportion of people eligible for any form of economic support that receive(d) it	HH, CS
111	Proportion of eligible people for legal services referred by a CHW to apply for them	CS
112	Proportion of eligible people that receive(d) legal services	HH, CS
41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH

Tuberculosis

127	Proportion of people confirmed with TB out of all people who were referred for diagnosis by CHW	CS
128	Proportion of people notified with TB through CHW referrals out of all people notified with TB	CS
129	Proportion of people to whom TB preventive treatment (TPT) support/follow-up was provided by CHW	CS
130	Proportion of people to whom TB disease treatment adherence support was provided	CS
131	TB contact screening	CS
132	Proportion of people who were successfully treated and who benefited from community-based TB treatment adherence support	CS
133	Proportion of individuals who started TPT who complete the course	CS

134	Proportion of people with TB referred by a CHW to a health facility to manage drug side effects	CS
109	Proportion of people eligible for any form of economic support referred by a CHW to apply for it	CS
110	Proportion of people eligible for any form of economic support that receive(d) it	HH, CS
111	Proportion of eligible people for legal services referred by a CHW to apply for them	CS
112	Proportion of eligible people that receive(d) legal services	HH, CS
41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH, CS

Neglected tropical diseases

60	Proportion of people receiving preventive chemotherapy for deworming	CS
135	Number of suspected neglected tropical disease (NTD) cases reported	HH, CS
136	Number of rumours of guinea worm disease/dracunculiasis cases reported	HH, CS
137	Number of rumoured yaws cases reported	HH, CS
138	Number of people referred to health centre for diagnosis or treatment of NTDs	HH, CS
139	Geographical coverage of preventive chemotherapy (PC) for targeted NTDs	HH
140	Population coverage of PC for targeted NTDs	HH
141	Proportion of households in the targeted communities that received social mobilization/awareness campaigns on NTDs	HH
142	Number of people bitten by an animal (animal bite cases) in the community, by animal	HH, CS
143	Number of deaths in the community occurring within three months after a snakebite or a dog bite	HH
144	Number of people screened for skin lesions consistent with NTDs (and population coverage)	HH, CS
145	Number of NTD cases that received adequate/recommended wound care	HH, CS
146	Number of people screened for signs and symptoms of visceral leishmaniasis (VL) and/or PKDL (and population coverage)	HH, CS
147	Proportion of people presenting hematuria, either visible hematuria reported by the patient or micro-hematuria detected by a positive dipstick	HH, CS
148	Proportion of people suffering from physical disability related to NTDs who receive rehabilitation support	HH, CS
149	Number (and proportion) of cases who received instructions for self-care for relevant NTDs	HH, CS
150	Proportion of targeted houses covered by domiciliary vector reduction measures	HH
151	Number of surface water bodies enumerated and mapped	HH
152	Proportion of households with all water storage containers covered and protected	HH
41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH, CS

Child protection and interpersonal violence

153	Number/proportion of households reached with health promotion messages about the impacts of violence against women and where to seek help	HH
154	Number of referrals made to health or other essential services for women and girls who disclose intimate partner violence or sexual violence	CS

34	Number/proportion of women aged 15–49 years old who have undergone female genital mutilation	HH
155	Number/proportion of girls under 15 years old who have undergone female genital mutilation (FGM) or are at risk of FGM	HH
156	Number/proportion of households with children where CHW raised awareness of positive parenting	HH
157	Number of households where CHW provided referrals to family services Module on Civil registration and vital statistics	CS

Civil registration and vital statistics

2	Number of live births	HH
45	Number of stillbirths (late fetal deaths)	HH
46	Number of neonatal deaths (0–27 days)	HH
55	Number of infant deaths (0 to less than 1 year old)	HH
56	Number of child deaths (1 to less than 5 years old)	HH
57	Number of child deaths (5 to less than 10 years old)	HH
82	Number of young adolescent deaths (10–14 years old)	HH
83	Number of older adolescent deaths (15–19 years old)	HH
158	Number of adult deaths	HH
44	Number of pregnancy-related deaths	HH
159	Number of deaths due to road traffic crashes	HH
160	Number of deaths due to drowning	HH
161	Number/proportion of live births happening in reporting period that were not registered, for which notification was submitted by CHW to local authorities	CS
162	Number/proportion of deaths happening in the reporting period that were not registered, for which notification was submitted by CHW to local authorities	CS
163	Number/proportion of children under-five year old whose births are registered with the civil authority	HH
164	Number/proportion of deaths that were registered in a timely manner over the reporting period	HH

Noncommunicable diseases

87	Proportion of adolescents currently using tobacco	HH, CS
165	Proportion of current tobacco users	HH, CS
80	Insufficient physical activity among children (0 to less than 5 years of age)	HH
81	Insufficient physical activity among children (5 to less than 10 years of age)	HH
86	Insufficient physical activity among adolescents	HH
166	Insufficient physical activity among adults	HH
68	Proportion of overweight among children under 5 years of age	HH, CS
69	Proportion of overweight and obesity in school-age children and adolescents 5–19 years	HH, CS
167	Proportion of overweight and obesity in adults	HH, CS
31	Proportion of women who have been screened for cervical cancer	HH
38	Proportion of people with raised blood pressure measured by CHW	HH, CS
168	Number/proportion of asymptomatic adults older than 40 with a BMI \geq 25 who have raised blood glucose/diabetes	HH, CS
89	Proportion of people with heavy episodic drinking	HH
169	Number/proportion of people supported for drug therapy and counselled to prevent heart attacks and stroke	CS
21	Proportion of households with primary reliance on clean fuels and technologies for cooking	HH

22	Proportion of households with primary reliance on clean fuels and technologies for lighting	HH
23	Proportion of households with primary reliance on clean fuels and technologies for heating	HH
49	Prevalence of low birthweight among newborns delivered in the community	CS

Nutrition

36	Proportion of antenatal care contacts during which women received breastfeeding counselling	CS
37	Proportion of community antenatal care contacts in the reporting period during which pregnant women were given/prescribed iron-containing supplements	CS
47	Proportion of newborns delivered in the community who were put to breast within the first hour of birth	CS
48	Proportion of newborns delivered in the community with documented birthweight	CS
49	Prevalence of low birthweight among newborns delivered in the community	CS
58	Proportion of consultations for infants under 6 months providing counselling on appropriate infant and young child feeding	CS
59	Proportion of consultations for children 6–23 months providing counselling on appropriate complementary feeding	CS
60	Proportion of people receiving preventive chemotherapy for deworming	CS
61	Percentage of children 6–59 months of age who received an age-appropriate dose of vitamin A through CHW contacts (routine contacts as well as contacts via events) in each semester	CS
62	Number/proportion of children aged 6–59 months assessed for wasting in the reporting period	CS
63	Proportion of children 6–59 months with mid-upper-arm circumference (MUAC) < 115 mm (severe acute malnutrition)	CS
64	Number/proportion of children under 5 years who had their weight assessed	CS
65	Number/proportion of children under 5 years who are underweight	HH, CS
66	Proportion of wasting among children aged 6–59 months	CS
67	Number/proportion of children under 5 years who had their height/length measured	CS
68	Proportion of overweight among children under 5 years of age	HH, CS
69	Proportion of overweight and obesity in school-age children and adolescents 5–19 years	HH, CS
70	Number/proportion of children who received treatment for diarrhoea	CS
84	Supplementation with iron-containing supplements (and folic acid) to adolescent girls and women of reproductive age for anaemia prevention	CS
85	Women and adolescent girls who are underweight	CS

Mental health

41	Number/proportion of people assessed for mental, neurological and substance use (MNS) disorders	CS
42	Number/proportion of people with mental, neurologic and substance use (MNS) referred	CS
43	Number/proportion of people with mental, neurologic and substance use (MNS) disorders receiving services	HH
92	Number of people with suicidal ideation or plan	CS
93	Number of people with suicide attempts	CS
94	Average time on weekdays and weekend days dedicated to screen time for leisure activities	HH
75	Proportion of children monitored for early signs of developmental delays	CS

76	Proportion of children with suspected developmental disabilities referred	CS
77	Proportion of caregivers being counselled on responsive caregiving and early learning activities	CS
78	Proportion of children with suspected developmental disorders and disabilities whose caregivers receive parenting information	CS
88	Proportion of alcohol consumers	HH
89	Proportion of people with heavy episodic drinking	HH
90	Proportion of people who use psychoactive drugs	HH
91	Proportion of people who inject psychoactive drugs	HH
170	Number/proportion of drug overdose deaths	HH, CS

People-centred services

109	Proportion of people eligible for any form of economic support referred by a CHW to apply for it	CS
110	Proportion of people eligible for any form of economic support that receive(d) it	HH, CS
111	Proportion of eligible people for legal services referred by a CHW to apply for them	CS
112	Proportion of eligible people that receive(d) legal services	HH, CS
171	Number of people who needed care and did not get it in the last month	HH
172	Proportion of people who refused care among those targeted by CHW	CS

Community-based surveillance /early warning

173	Event/alert case detected	CS
174	Proportion of CBS alerts responded to within 24 hours or within specified time period from the CBS protocol	CS
175	Proportion of communities in which action was taken following an alert (per month)	CS





Unicef

Division of Communication

3 United Nations Plaza

New York, NY 10017, USA

www.unicef.org